



|            |                                                                                               |
|------------|-----------------------------------------------------------------------------------------------|
| Title      | Auslander-Reiten quivers of orders Dedicated to Professor Hisao Tominaga on his 60th birthday |
| Author(s)  | Nishida, Kenji                                                                                |
| Citation   | 長崎大学教養部紀要. 自然科学篇. 1987, 28(1), p.1-9                                                          |
| Issue Date | 1987-07                                                                                       |
| URL        | <a href="http://hdl.handle.net/10069/16585">http://hdl.handle.net/10069/16585</a>             |
| Right      |                                                                                               |

This document is downloaded at: 2019-04-24T20:17:59Z

## Auslander-Reiten quivers of orders

Dedicated to Professor Hisao Tominaga on his 60th birthday

Kenji NISHIDA

(Received March 16, 1987)

### 1. Introduction

Let  $R$  be a complete discrete valuation ring with prime element  $\pi$  and residue field  $k$ . Let  $A$  be an  $R$ -order in the semisimple finite dimensional algebra over the quotient field of  $R$  and  $\Gamma$  a hereditary  $R$ -order in the same algebra such that  $J = \text{rad } \Gamma \subset A \subset \Gamma$ . Put  $A = A/J$  and  $B = \Gamma/J$ . Then  $A$  is a  $k$ -subalgebra of the semisimple  $k$ -algebra  $B$ . We assume that  $A$  is basic, so that  $B$  is, too. By the technical reason we assume that  $A$  is not a hereditary order whose structure is completely determined (see, for example [4]). In the previous paper [3], we showed that  $\text{latt } A$ , the category of all right  $A$ -lattices, is representation equivalent to  $\text{mod}_{\text{sp}} C$  with  $C = \begin{pmatrix} A & DI \\ 0 & K \end{pmatrix}$  under the above situation. But, as for the Auslander-Reiten quiver of  $\text{latt } A$ , we didn't give any information there, while Roggenkamp[8] gives the method of constructing it under the further assumption that  $\text{rad } A$  decomposes into a projective  $A$ -lattice and a  $\Gamma$ -lattice and  $A$  is of finite lattice type. In fact, for  $C' = \begin{pmatrix} A & B \\ 0 & B \end{pmatrix}$ , the relation of irreducible maps between  $\text{latt } A$  and the subcategory  $\mathcal{C}$  of  $\text{mod } C'$  is determined in [8]. It is noted that  $C'$  is a hereditary algebra under the assumption of [8]. When  $A$  is of finite lattice type, the Auslander-Reiten quiver of  $\text{latt } A$  is obtained from that of  $\mathcal{C}$  by identifying some vertices of  $\mathcal{C}$  [8, III Theorem V].

In this paper, we generalize the results of [8] and study the problem how the Auslander-Reiten quiver of  $\text{latt } A$  is constructed from that of  $\text{mod}_{\text{sp}} C$  under our situation. In order to describe our results more precisely we prepare notation. We identify a  $C'$ -module with a triple  $(X, Y, \phi)$  where  $X$  is a right  $A$ -module,  $Y$  a right  $B$ -module and  $\phi : X \otimes_A B \rightarrow Y$  a  $B$ -homomorphism. Let  $\mathcal{C}$  be a full subcategory of  $\text{mod } C'$  consisting of the modules of the form  $(X, Y, \phi)$  such that  $X$  is finitely generated, the adjoint  $\bar{\phi} \in \text{Hom}_A(X, Y)$  of  $\phi$  is injective and  $(\text{Im } \phi)B = Y$ . For a  $A$ -lattice  $M$ , put  $\bar{M} = M/MJ$ ,

$\overline{M\Gamma} = M\Gamma/MJ$  and  $\bar{\phi} : \overline{M} \rightarrow \overline{M\Gamma}$  the canonical inclusion. Since  $\bar{\phi}$  induces  $\phi' : \overline{M} \otimes_A B \rightarrow \overline{M\Gamma}$ , we may put  $H(M) = (\overline{M}, \overline{M\Gamma}, \phi')$ . We sometimes omit  $\phi'$  and write  $H(M) = (\overline{M}, \overline{M\Gamma})$ . Then  $H$  is a functor from  $\text{latt } A$  to  $\text{mod } C'$  and the following was proved in [2, 5].

**THEOREM A.** *There exists a representation equivalence  $\text{latt } A \approx \mathcal{C}$  induced from  $H$ .*

A  $k$ -algebra  $Z$  is called a right peak ring, if the socle  $\text{soc } Z$  of  $Z$  as a right  $Z$ -module is projective. A right peak ring is introduced and studied in [9] when  $\text{soc } Z$  is homogeneous, however, almost all results about a right peak ring with a homogeneous socle hold for nonhomogeneous case. We next define a right peak  $k$ -algebra which plays a valuable part in this paper. Let  $G_1, \dots, G_t$  be the representatives of nonisomorphic indecomposable projective right  $\Gamma$ -lattices. Put  $S_i = G_i/G_iJ$  ( $i=1, \dots, t$ ) and  $I = S_1 \oplus \dots \oplus S_t$ . Then  $S_1, \dots, S_t$  are the representatives of nonisomorphic simple right  $B$ -modules. For  $K_i = \text{End}_B S_i$  ( $i=1, \dots, t$ ) put  $K = K_1 \otimes \dots \otimes K_t$  a product of division rings. Let  $D$  be the usual duality  $D(-) = \text{Hom}_k(-, k)$ . Then it is easily seen that a ring  $C = \begin{pmatrix} A & DI \\ 0 & K \end{pmatrix}$  is a right peak ring by [9]. For  $u = (X, Y, \psi) \in \text{mod } C'$ , put  $H'(u) = (X, Y \otimes_B DI, \bar{\psi})$ , where  $\bar{\psi} : X \otimes_A DI \rightarrow Y \otimes_B DI$  is given by  $\bar{\psi}(x \otimes g) = \psi(x \otimes 1) \otimes g$  ( $x \in X, g \in DI$ ). Then  $H'$  is a functor from  $\text{mod } C'$  to  $\text{mod } C$  which is a category equivalence. Let  $\text{mod}_{\text{sp}} C$  be the full subcategory of  $\text{mod } C$  whose modules have projective socles and  $\text{mod}_{\text{sp}}^1 C$  the full subcategory of  $\text{mod}_{\text{sp}} C$  consisting of all modules having no direct summand of the form  $(0, K_i, 0)$  ( $i=1, \dots, t$ ), that is, having no simple projective direct summand. Then the following was proved in [3, Theorem 4].

**THEOREM B.** *There exists a representation equivalence  $\text{latt } A \approx \text{mod}_{\text{sp}}^1 C$  induced from  $\Phi = H'H$ .*

Since  $\text{mod}_{\text{sp}} C$  has been extensively studied by Simson[9] when  $C$  is a right peak ring, the well understanding of the functor  $\Phi$  contributes the investigation of the representation theory of the orders. We consider the behavior of irreducible maps, almost split sequences under the functor  $\Phi$  and then provide the method of constructing the Auslander-Reiten quiver of  $\text{latt } A$  from that of  $\text{mod}_{\text{sp}} C$ . Hence our main theorem is the following, where the undefined notation will be explained in the next section.

**THEOREM C.** *The Auslander-Reiten quiver of latt  $\Lambda$  is obtained from that of  $\text{mod}_{\text{sp}} C$  by identifying the indecomposable injective  $C$ -module  $\Phi(G)$  with the simple projective  $C$ -module  $(0, \text{End}_B \overline{\theta(G)}, 0)$  for every indecomposable projective  $\Gamma$ -lattice  $G$ .*

We assume throughout the paper that all modules are finitely generated. The reader is referred to [1, 8] for the definition and the properties of irreducible maps, almost split sequences, Auslander-Reiten quivers and to [9] for properties of  $\text{mod}_{\text{sp}} C$  for a right peak ring  $C$ , for example, it was proved in [9] that  $\text{mod}_{\text{sp}} C$  has almost split sequences, enough injectives, and so on.

## 2. Auslander-Reiten quivers of latt $\Lambda$

The functor  $\Phi$  is essentially dominated by the functor  $H$ . The useful results about  $H$  is provided in [5, §1], so we will freely use them. Firstly we begin with a lemma which seems to be well-known, but we provide the proof here for the completeness.

**LEMMA 1.** *For  $M, M' \in \text{latt } \Lambda$ ,  $f: M \rightarrow M'$  is a splitting monomorphism (respectively epimorphism) if and only if  $\Phi(f): \Phi(M) \rightarrow \Phi(M')$  is a splitting monomorphism (respectively epimorphism).*

**PROOF.** Since  $H$  is a category equivalence, we prove the lemma for  $H$ . Let  $H(f)$  be a splitting monomorphism. Then there exists  $g: M' \rightarrow M$  such that  $H(g)H(f) = 1_{H(M)}$ . Define  $f_1: M\Gamma \rightarrow M'\Gamma$  and  $g_1: M'\Gamma \rightarrow M\Gamma$  by  $f_1(m\gamma) = f(m)\gamma$  and  $g_1(m'\gamma) = g(m')\gamma$  for  $m \in M, m' \in M', \gamma \in \Gamma$ . By assumption it holds that  $g_1 f_1(M\Gamma) + MJ = M$ , so that  $g_1 f_1(M\Gamma) = M$  by Nakayama's Lemma. Since  $\text{Ker } g_1 f_1 = 0$  by the rank argument,  $g_1 f_1$  is an isomorphism. On  $M$   $g_1 f_1$  equals  $gf$ , thus  $gf$  is also an isomorphism. Therefore,  $f$  is a splitting monomorphism. The rest of the proof is almost trivial.

The following two lemmas are also obtained in [8, Proposition 4.4] with a slightly different manner.

**LEMMA 2.** *Let  $M, N \in \text{latt } \Lambda$  with  $M$  indecomposable and let  $f: M \rightarrow N$  be an irreducible map. Then  $\Phi(f) = 0$  if and only if  $f(M)$  is a direct summand of  $NJ$ . Moreover, in this case  $N$  is a projective  $\Lambda$ -lattice.*

**PROOF.** It holds that  $\Phi(f) = 0 \Leftrightarrow H(f) = 0 \Leftrightarrow f(M) \subset NJ$ . Thus if  $\Phi(f) = 0$ ,

then  $f$  factors through  $M \xrightarrow{f} NJ \subset N$ . Since  $f$  is irreducible,  $f: M \rightarrow NJ$  is a splitting monomorphism. The converse is obvious. The rest of the statement is showed in [8, Proposition 4. 4].

LEMMA 3. *Let  $M, N$  be as in Lemma 2. Assume that  $\Phi(f)$  is a nonzero nonisomorphism for  $f: M \rightarrow N$ . Then  $f$  is irreducible in  $\text{latt } \Lambda$  if and only if  $\Phi(f)$  is irreducible in  $\text{mod}_{\text{sp}} C$ .*

PROOF. Let  $f_1: M\Gamma \rightarrow N\Gamma$  be as in the proof of Lemma 1, and let  $\bar{f}: \bar{M} \rightarrow \bar{N}$ , resp.  $\bar{f}_1: \bar{M}\Gamma \rightarrow \bar{N}\Gamma$  be induced from  $f$ , resp.  $f_1$ . Assume that  $\Phi(f)$  factors through  $\Phi(M) \xrightarrow{g} u \xrightarrow{h} \Phi(N)$ . Since only simple projective  $C$ -modules are not in  $\text{Im } \Phi$  among the modules in  $\text{mod}_{\text{sp}} C$  by Theorem B, we can assume that  $u \in \text{Im } \Phi$ , so that  $u = \Phi(X)$ ,  $g = \Phi(\alpha)$ ,  $h = \Phi(\beta)$  for  $X \in \text{latt } \Lambda$ ,  $\alpha: M \rightarrow X$ ,  $\beta: X \rightarrow N$ . Put  $\phi = f - \beta\alpha: M \rightarrow N$ . Then  $\text{Im } \phi \subset NJ$  by  $\bar{f} = \bar{\beta}\bar{\alpha}$ . Hence  $f = \phi + \beta\alpha = (\iota, \beta)(\phi, \alpha)$ , where  $\iota: NJ \rightarrow N$  is a canonical inclusion. If  $f$  is irreducible, then either  $(\iota, \beta)$  is a splitting epimorphism or  $(\phi, \alpha)$  is a splitting monomorphism. Assume that  $(\iota, \beta)$  is a splitting epimorphism. Then there exists  $(\gamma, \beta'): N \rightarrow NJ \oplus X$  such that  $\beta\beta' + \iota\gamma = 1_N$ . It holds that  $\text{Im } \beta\beta' + NJ = N$  and so  $\text{Im } \beta\beta' = N$ . Therefore,  $\beta\beta': N \rightarrow N$  is an isomorphism. Hence we have that  $\beta$  is a splitting epimorphism, and  $H(\beta)$  is, too. Similarly, if  $(\phi, \alpha)$  is a splitting monomorphism, then  $H(\alpha)$  is a splitting monomorphism. Hence we showed that  $\Phi(f)$  is irreducible. The converse is obvious by Lemma 1.

PROPOSITION 1. *For every  $\Gamma$ -lattice  $M$ ,  $\Phi(M)_C$  is injective.*

PROOF. We assume that  $M = G_i$  for some  $i(1 \leq i \leq t)$ . Thus  $H(G_i) = (S_i, S_i, \phi_i)$  with  $\phi_i: S_i \otimes_A B \rightarrow S_i$  canonically, and  $\Phi(G_i) = (S_i, S_i \otimes_B DI, \psi_i)$ . Since  $S_i \otimes_B DI \cong K_i$  and  $S_i \otimes_A DI \cong S_i \otimes_A I^*$  with  $I^* = \text{Hom}_K(I, K)$  we conclude that the  $K_i$ -homomorphism  $\psi_i: S_i \otimes_A DI \rightarrow K_i$  is the canonical evaluation map and  $\Phi(G_i) \cong (S_i, K_i, \psi_i)$ . Then  $D\Phi(G_i) \cong ({}_A S_i^*, {}_K K_i, \lambda_i)$ , where  $\lambda_i: I^* \otimes_K K_i \rightarrow S_i^*$  is given by  $[(f_j) \otimes y \rightarrow f_i y]$  for  $f_j \in S_j^*$  ( $j=1, \dots, t$ ), is a projective left  $C$ -module by the form of  $C$ . Therefore,  $\Phi(G_i)_C$  is injective.

COROLLARY 1. *Let  $f: M \rightarrow N$  be an irreducible map between indecomposable  $\Lambda$ -lattices such that  $M$  is a  $\Gamma$ -lattice and  $\Phi(f) \neq 0$ . Then  $M$  and  $N$  are isomorphic  $\Lambda$ -lattices.*

PROOF. If  $\Phi(f)$  is not an isomorphism, then  $\Phi(f)$  is irreducible by Lemma 3. Since  $\Phi(M)$  is injective by Proposition 1,  $\Phi(f)$  is an epimorphism. However,  $\Phi(M)$  has a projective simple socle and  $\Phi(N) \in \text{mod}_{\text{sp}} C$ , a contradiction. Thus  $\Phi(f)$  is an isomorphism and  $M \cong N$ .

Since the Auslander-Reiten quiver of an order such that there exists an irreducible map  $M \rightarrow M$  is completely determined by [11] (cf. also [7, Theorem 3.11]), we assume in the following that there exists no irreducible map  $M \rightarrow M$  for every  $A$ -lattice  $M$ .

**COROLLARY 2.** *Let  $f: M \rightarrow N$  be an irreducible map between  $A$ -lattices. Then  $\Phi(f) \neq 0$  if and only if  $M$  is not a  $\Gamma$ -lattice.*

The next proposition is essential, because it assures that the almost split sequences, except those starting from a  $\Gamma$ -lattice, are preserved by  $\Phi$ .

**PROPOSITION 2.** *Let  $0 \rightarrow M \xrightarrow{f} M' \xrightarrow{g} M'' \rightarrow 0$  be an exact sequence of non-zero  $A$ -lattices such that  $f$  is irreducible and  $M$  is indecomposable and not a  $\Gamma$ -lattice. Then  $0 \rightarrow \Phi(M) \xrightarrow{\Phi(f)} \Phi(M') \xrightarrow{\Phi(g)} \Phi(M'') \rightarrow 0$  is exact.*

**PROOF.** Let  $f_1$  and  $g_1$  be the same as in the proof of Lemma 1. Then we have the following two exact sequences,

$$\begin{aligned} 0 \rightarrow \ker g_1 \rightarrow M'\Gamma \xrightarrow{g_1} M''\Gamma \rightarrow 0, \\ 0 \rightarrow M\Gamma \xrightarrow{f_1} \ker g_1 \rightarrow T \rightarrow 0, \end{aligned}$$

where  $T$  is a torsion  $R$ -module. Since  $H(f) \neq 0$  by Corollary 2, it holds that  $\text{Im } f_1 \not\subset (\ker g_1)J$ . Thus we have a decomposition  $\ker g_1 = X_1 \oplus X_2$ , where  $X_1, X_2$  are  $\Gamma$ -lattices with  $X_1 \neq 0$  such that  $f_1 = (f'_1, f'_2)$  for  $f'_i: M\Gamma \rightarrow X_i (i=1, 2)$ ,  $\text{Im } f'_i = X_1$  and  $\text{Im } f'_2 \subset X_2J$ . Considering the induced map  $\bar{f}_1 = (\bar{f}'_1, \bar{f}'_2): \overline{M\Gamma} \rightarrow \overline{X_1 \oplus X_2} = \overline{\ker g_1}$ , we have  $\bar{f}'_2 = 0$  and  $\bar{f}'_1$  is surjective. Put  $T_1 = \ker \bar{f}'_1$ . Then  $\ker H(f) = (0, T_1, 0)$  by [6, Lemma 1.4]. On the other hand, since  $H(f)$  is irreducible,  $H(f)$  is a proper epimorphism or proper monomorphism. It cannot be an epimorphism, so a monomorphism. It holds that  $T_1 = 0$  and then  $\bar{f}'_1$  is an isomorphism. Thus we have  $M\Gamma \cong X_1$ . Since  $\text{rank}_R M\Gamma = \text{rank}_R (\ker g_1) = \text{rank}_R X_1 + \text{rank}_R X_2$ , we conclude that  $X_2 = 0$  and  $\text{Im } f_1 = \ker g_1$ , so that  $0 \rightarrow M\Gamma \rightarrow M'\Gamma \rightarrow M''\Gamma \rightarrow 0$  is exact. Hence by [6, Lemma 1.4]  $0 \rightarrow \Phi(M) \rightarrow \Phi(M') \rightarrow \Phi(M'') \rightarrow 0$  is exact.

**COROLLARY 3.** *Let  $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$  be an almost split sequence in latt  $A$  such that  $M$  is not a  $\Gamma$ -lattice. Then  $0 \rightarrow \Phi(M) \rightarrow \Phi(M') \rightarrow \Phi(M'') \rightarrow 0$  is an almost split sequence in  $\text{mod}_{\text{sp}} C$ .*

**PROOF.** By Proposition 2  $0 \rightarrow \Phi(M) \rightarrow \Phi(M') \rightarrow \Phi(M'') \rightarrow 0$  is exact and by Lemma 1 it doesn't split. Let  $0 \rightarrow \Phi(M) \rightarrow \Phi(N) \rightarrow L \rightarrow 0$  be an almost split sequence starting from  $\Phi(M)$  in  $\text{mod}_{\text{sp}} C$ . Then there exists a splitting

epimorphism  $\Phi(N) \rightarrow \Phi(M')$ . We have a splitting epimorphism  $N \rightarrow M'$  by Lemma 1. Applying the similar argument to the irreducible map  $M \rightarrow N$  and the almost split sequence  $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$  we have a splitting epimorphism  $M' \rightarrow N$ . Therefore,  $M' \cong N$ .

By Corollary 3 the remaining almost split sequences of latt  $\Lambda$  are those starting from  $\Gamma$ -lattices. Before investigating these almost split sequences we summarize the projective  $C$ -modules. An indecomposable projective  $C$ -module, up to isomorphism, is either a simple projective module of the form  $(0, K_i, 0)$  ( $i=1, \dots, t$ ) or  $\Phi(M)$  for an indecomposable projective  $\Lambda$ -lattice  $M$ . For it holds that a  $C$ -module  $X$  is projective if and only if  $X$  is a projective object in  $\text{mod}_{\text{sp}} C$ , since  $C$  is a right peak ring. Thus statement follows from Theorem B.

In order to describe the behavior of the almost split sequences starting from  $\Gamma$ -lattices under the functor  $\Phi$ , we prepare some well-known results about hereditary orders [4, Chapter 39]. Decompose  $\Gamma = \Gamma_1 \oplus \dots \oplus \Gamma_k$  with each  $\Gamma_i$  indecomposable as a ring. Let  $G_{ij}$  ( $j=1, \dots, a_i$ ) be the representatives of the indecomposable projective  $\Gamma_i$ -lattices ( $i=1, \dots, k$ ). We number as follows for each  $i$  ( $i=1, \dots, k$ ) [4, (39.8)];  $G_{ij} \cong G_{i, j+1}(\text{rad } \Gamma_i)$  ( $j=i, \dots, a_i - 1$ ) and  $G_{ia_i} \cong G_{i1}(\text{rad } \Gamma_i)$ . Put

$$\theta(G_{ij}) = \begin{cases} G_{i, j+1}, & \text{if } j=1, \dots, a_i - 1 \\ G_{ij}, & \text{if } j=a_i. \end{cases}$$

Then  $\theta(G_{ij})$  is the unique minimal over  $\Gamma$ -module of  $G_{ij}$  such that  $\theta(G_{ij})/G_{ij}$  is a simple  $\Gamma$ -module. Of course, in the case that  $j=a_i$  we identify  $G_{ia_i}$  to its isomorphic image in  $G_{i1}$ .

**PROPOSITION 3.** *Let  $G$  be an indecomposable  $\Gamma$ -lattice. Then the following holds.*

1) *If there exists an almost split sequence  $0 \rightarrow G \xrightarrow{f} L \xrightarrow{g} M \rightarrow 0$  in latt  $\Lambda$ , then  $0 \rightarrow p \rightarrow \Phi(L) \rightarrow \Phi(M) \rightarrow 0$  is an almost split sequence in  $\text{mod}_{\text{sp}} C$  and  $p \cong (0, F, 0)$  is a simple projective  $C$ -module where  $F = \text{End}_{\text{sp}} \overline{(G)}$  is a division ring.*

2) *If  $G$  is an injective  $\Lambda$ -lattice and  $f: G \rightarrow N$  is an irreducible map in latt  $\Lambda$ , then there exists an irreducible map  $(0, F, 0) \rightarrow \Phi(N)$ , where  $F$  is the same as in 1.*

**PROOF.** 1) Since  $\Phi(M)$  is not projective and  $\Phi(g)$  is irreducible, the almost split sequence ending at  $\Phi(M)$  in  $\text{mod}_{\text{sp}} C$  is  $0 \rightarrow p \rightarrow \Phi(L) \rightarrow \Phi(M) \rightarrow 0$ . In order to prove  $p \cong (0, F, 0)$  it suffices to show  $(0, F, 0) \cong \ker \Phi(g)$ . Let

$g_1$  be the same as in the proof of Lemma 1. Then  $g_1$  is a splitting epimorphism and  $L\Gamma = \ker g_1 \oplus Y$ . Decompose  $L\Gamma$  into a direct sum of indecomposable projective  $\Gamma$ -lattices, say  $L\Gamma = L_1 \oplus \dots \oplus L_m$ . Then  $LJ = L_1J \oplus \dots \oplus L_mJ$  is also a direct sum of indecomposable  $\Gamma$ -lattices. Since  $f(G)$  is a direct summand of  $LJ$  by Lemma 2, we can assume  $f(G) = L_1J$  and  $\theta(G) = L_1$ . It holds that  $L\Gamma = \theta(G) \oplus X$  for a  $\Gamma$ -lattice  $X$ . Since  $g_1f(G) = 0$  and  $\theta(G)/G$  is  $R$ -torsion, we conclude that  $g_1(\theta(G)) = 0$ . By rank argument we have that  $\ker g_1 \cap X = 0$ . Then by an elementwise argument  $\ker g_1 \oplus Y = \theta(G) \oplus X$  deduces that  $\ker g_1 = \theta(G)$ , so that  $\ker \bar{g}_1 = \overline{\theta(G)}$  for  $\bar{g}_1: \overline{L\Gamma} \rightarrow \overline{MJ}$ . On the other hand, since  $\text{Im } f \subset LJ$ ,  $\bar{g}: \overline{L} \rightarrow \overline{M}$  is an isomorphism. It holds that  $\ker H(g) \cong (0, \overline{\theta(G)}, 0)$ . Applying the functor  $H'$  we have that  $\ker \Phi(g) \cong (0, F, 0)$ .

2) By Corollary 2  $\Phi(f) = 0$  and by Lemma 2  $\text{Im } f$  is a direct summand of  $NJ$ . Thus the irreducible map  $f$  is a proper monomorphism such that  $\text{coker } f$  is a simple  $A$ -module by [10, Proposition 1.2], since  $G$  is an injective  $A$ -lattice. It holds that  $\text{Im } f = NJ$  and  $NJ = \text{rad}_A N$ , so  $N\Gamma$  is an indecomposable  $\Gamma$ -lattice. Thus we conclude that  $N\Gamma \cong \theta(G)$ , so that  $\text{rad } H(N) = \text{rad } (\overline{N}, \overline{N\Gamma}) \cong (0, \overline{\theta(G)}, 0)$ . Applying the functor  $H'$  it holds that  $\text{rad } \Phi(N) \cong (0, F, 0)$ . By Lemma 2 and the paragraph succeeding Corollary 3,  $\Phi(N)$  is projective. Hence there exists an irreducible map  $(0, F, 0) \rightarrow \Phi(N)$ .

**COROLLARY 4.** *If  $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$  is an exact sequence in latt  $A$  such that  $0 \rightarrow \Phi(M) \rightarrow \Phi(M') \rightarrow \Phi(M'') \rightarrow 0$  is an almost split sequence in  $\text{mod}_{\text{sp}} C$ . then it is an almost split sequence in latt  $A$ .*

**PROOF.** Keeping in mind that  $M''$  is not a projective  $A$ -lattice this follows from Proposition 3 and Corollary 3.

**PROOF OF THEOREM C.** This follows directly from Lemma 3, Proposition 3 and Corollaries 3, 4.

Following the notation provided before Proposition 3, we restate Theorem C more concretely. Put  $p_{ij} = (0, \text{End}_B \overline{G}_{ij}, 0)$  ( $i = 1, \dots, k$ ;  $j = 1, \dots, a_i$ ). Then they are simple projective  $C$ -modules and each simple projective  $C$ -module is isomorphic to one of them. Note that  $E(p_{ij})$ , the injective hull of  $p_{ij}$ , is isomorphic to  $\Phi(G_{ij})$  for all  $i, j$ . Let  $Q$  be the Auslander-Reiten quiver of  $\text{mod}_{\text{sp}} C$ . Then  $Q$  contains each  $p_{ij}$  and  $E(p_{ij})$  as its vertices. In  $Q$ , identify  $E(p_{ij})$  with  $p_{i, j+1}$  ( $j = 1, \dots, a_i - 1$ ) and  $E(p_{ia_i})$  with  $p_{i1}$  for all  $i = 1, \dots, k$ , and then put the quiver obtained after this identification  $Q'$ . Then  $Q'$  is the Auslander-Reiten quiver of latt  $A$ , where the vertices identified by the above

way correspond each other to indecomposable  $\Gamma$ -lattices. More precisely,  $E(p_{ij}), p_{i, j+1}$  in  $Q$  correspond to  $G_{ij}$  in  $Q'$  ( $j=1, \dots, a_i - 1$ ) and  $E(p_{ia_i}), p_{i1}$  in  $Q$  to  $G_{ia_i}$  in  $Q'$  for all  $i$ .


EXAMPLE. Let  $A = \begin{pmatrix} R & R & R & R & R \\ r & R & r & R & R \\ r & r & R & R & r \\ r & r & r & R & r \\ r & r & r & r & R \end{pmatrix}, \Gamma_1 = (R)_5, \Gamma_2 = \begin{pmatrix} R & R \dots R \\ r & R \dots R \\ \vdots & \vdots \quad \vdots \\ \vdots & \vdots \quad \vdots \\ r & R \dots R \end{pmatrix}$

with  $r = \pi R$ . Then  $A \subset \Gamma_1$  and  $A \subset \Gamma_2$  satisfy our assumption. Firstly we consider the case of  $\Gamma_1$ . Since  $\text{rad } \Gamma_1 = (r)_5$ , we have


$$C = \begin{pmatrix} k & k & k & k & k & k \\ 0 & k & 0 & k & k & k \\ 0 & 0 & k & k & 0 & k \\ 0 & 0 & 0 & k & 0 & k \\ 0 & 0 & 0 & 0 & k & k \\ 0 & 0 & 0 & 0 & 0 & k \end{pmatrix}$$

which is a path algebra of the bounden quiver  $\begin{matrix} \circ \rightarrow \circ \rightarrow \circ \\ \downarrow \quad \downarrow \quad \downarrow \\ \circ \rightarrow \circ \rightarrow \circ \end{matrix}$  with commuting cycles.

The Auslander-Reiten quiver of  $\text{mod}_{\text{sp}} C$  is the following ;




where  $p$  is simple projective and  $E(p)$  its injective hull and the dotted lines indicate the  $\tau$ -orbit of the Auslander-Reiten translation  $\tau$ . Identifying  $p$  with  $E(p)$  we get the Auslander-Reiten quiver of  $A$  ;



where  $G_2$  is a  $\Gamma_1$ -lattice. For the case of  $\Gamma_2$ ,  $C$  is a path algebra of the bounden quiver  $\begin{matrix} \circ \rightarrow \circ \\ \downarrow \quad \downarrow \\ \circ \rightarrow \circ \end{matrix}$  with a commuting cycle. The Auslander-

Reiten quiver of  $\text{mod}_{\text{sp}} C$  is the following ;



where  $p_1$  and  $E(p_i)$  ( $i=1, 2$ ) are the same as in the first case. Identifying  $p_1$  with  $E(p_2)$  and  $p_2$  with  $E(p_1)$  we get the same Auslander-Reiten quiver of latt  $A$ . In this case,  $G_1 = (rRRRR)$  and  $G_2 = (RRRRR)$  are  $\Gamma_2$ -lattices.

### References

- [ 1 ] Auslander, M., Reiten, I. : Representation theory of Artin algebras III, IV, Comm. Algebra, **3**(1975), 239–294, *ibid.*, **5**(1977), 443–518.
- [ 2 ] Green, E. L., Reiner, I. : Integral representations and diagrams, Michigan Math. J., **25**(1978), 53–84.
- [ 3 ] Nishida, K. : Representations of orders and vector space categories, J. P. A. Algebra **33**(1984), 209–217.
- [ 4 ] Reiner, I. : Maximal Orders (Academic Press, New York, 1975).
- [ 5 ] Ringel, C. M., Roggenkamp, K. W. : Diagrammatic methods in the representation theories of orders, J. Algebra **60**(1979), 11–42.
- [ 6 ] Roggenkamp, K. W. : Orders of global dimension two, Math. Z., **160**(1978), 63–67.
- [ 7 ] Roggenkamp, K. W. : The lattice type of orders II, Integral Representations and applications (L. N. M. **882**, Springer, Berlin, 1981), 430–477.
- [ 8 ] Roggenkamp, K. W. : Auslander-Reiten species for socle determined categories of hereditary algebras and for generalized Backstrom orders (Mitt. Math. Seminar, Giessen, **159**, 1983)
- [ 9 ] Simson, D. : Vector space categories, right peak rings and their socle projective modules, J. Algebra, **92**(1985), 532–571.
- [10] Schmidt, J. W. : Irreducible homomorphisms for lattices over orders, Bull. Austral. Math. Soc., **17**(1977), 109–124.
- [11] Wiedemann, A. : Orders with loops in their Auslander-Reiten graph, Comm. Algebra, **9**(1981), 641–656.