



Title	Effect of in Vivo Administration of Anti-Lyt-2 mAb and Anti-L3T4 mAb on Tumor Rejection
Author(s)	Nakayama, Eiichi; Mieno, Masahiro; Shiku, Hiroshi
Citation	Acta Medica Nagasakiensia. 1989, 34(1), p.3-9
Issue Date	1989-03
URL	http://hdl.handle.net/10069/17526
Right	

This document is downloaded at: 2019-04-24T16:51:28Z

Effect of in Vivo Administration of Anti-Lyt-2 mAb and Anti-L3T4 mAb on Tumor Rejection

Eiichi Nakayama, Masahiro Mieno and Hiroshi Shiku

*Department of Oncology,
Nagasaki University School of Medicine*

INTRODUCTION

Effector cells mediating tumor rejection were controversial. Several investigators have shown that by utilizing adoptive transfer into adult, thymectomized, bone marrow reconstituted (ATXBM) or athymic mice, purified Lyt-2⁺ cells or cloned Lyt-2⁺ cytotoxic T-cells were capable for tumor rejection (1,6). On the other hand, Lyt-2⁻ (L3T4⁺) cells have also been shown to mediate tumor rejection (7,8). These studies suggested that either Lyt-2⁺ or L3T4⁺ cells appeared to be capable of mediating the rejection. However, it is still unknown whether either type of T-cells is predominantly involved or either Lyt-2⁺ or L3T4⁺ cells themselves are fully capable of mediating the rejection. In this study, by eliminating Lyt-2⁺ cells and/or L3T4⁺ cells in vivo by administrating anti-Lyt-2.2 mAb, anti-L3T4 mAb or both, we directly investigated the T-cell type that was involved in the effector mechanisms of tumor rejections.

MATERIALS AND METHODS

Mice. C57BL/6 (B6), BALB/c, and (BALB/c x C57BL/6) F₁ (CB6F₁) mice were purchased from Shizuoka Laboratory Animal Center (Shizuoka, Japan). B6-Lyt-1.1 and B6-Lyt-2.1, 3.1 mice were originally provided by Dr. E.A. Boyse, Memorial Sloan-Kettering Cancer Center, New York and bred in our laboratory. The Lyt phenotypes of B6, B6 Lyt congenic, and

BALB/c mice are as follows : B6, Lyt-1.2,2.2, 3.2 ; B6-Lyt-1.1, Lyt-1.1,2.2,3.2 ; B6-Lyt-2.1,3.1, Lyt-1.2,2.1,3.1 ; BALB/c, Lyt-1.2,2.2,3.2.

Tumors. B6RV 2 and BALBRVD leukemias induced by injection of neonatal B6 and BALB/c mice respectively, with radiation leukemia virus (9). RL♂1 is a radiation-induced leukemia of BALB/c origin (10,11). These tumors were maintained in ascites form in the strain of origin.

Monoclonal Antibodies. Anti-L3T4 mAb, a rat antibody of the IgG 2b immunoglobulin class, produced by hybridoma GK1.5 (12,13), was kindly provided by Dr. F. Fitch, University of Chicago (Chicago, IL). Anti-Lyt-2.2 mAb and other mAb used have been described (14). The titers of both anti-L3T4 and anti-Lyt-2.2 mAb determined by antibody-mediated complement-dependent cytotoxicity assay were 1 : 20,000. These antibodies were used in the form of ascites from hybridoma bearing mice. The concentrations of anti-L3T4 and anti-Lyt-2.2 mAb in pooled ascites were 2.8 and 7.1 mg/ml, respectively, as quantified by protein assay (Bio-Rad Laboratories, Richmond, CA), and by quantitative cellulose acetate electrophoresis.

Tumor Assay. Tumors were harvested in Eagles' minimum essential medium (MEM) and washed twice with medium. Then the desired number of tumor cells (in 0.2 ml) was injected intradermally into the backs of mice through a 30-gauge stainless steel needle. Before inoculation of tumor cells, the hair was shaved with

Abbreviations used : ATXBM, adult thymectomized, bone marrow reconstituted ; B6, C57BL/6 ; CB6F₁, (BALB/c x C57BL/6) F₁ ; MEM, minimum essential medium

clippers. The diameters of tumors were measured with vernier calipers, twice, at right angles, to calculate the mean diameter.

Intravenous Injection of Antibody. Mice were anesthetized with ether and injected through the retrobulbar venous plexus with 0.2 ml of antibodies (ascites), diluted 1 : 4 with MEM, through a 1-ml disposable tuberculin syringe fitted with a 26-gauge needle (Terumo, Inc., Tokyo).

Antibody-mediated, Complement-Dependent Cytotoxicity. Tests were performed as described previously (15). Before tests, dead cells were removed by density gradient centrifugation in bovine serum albumin (16).

RESULTS

Clearance of anti-Lyt-2.2 mAb and anti-L3T4 mAb after in vivo administration. Clearance of anti-Lyt-2.2 and anti-L3T4 mAb in mice was studied by the cytotoxic test with B6 thymocytes as target cells, using serum specimens obtained every other day after in vivo administration of mAb. As shown in Fig. 1, anti-Lyt-2.2 mAb remained until day 30 and disappeared around day 33-35. On the other hand, anti-L3T4 mAb disappeared around day 10. Because of rather rapid clearance of L3T4 mAb, we injected L3T4 mAb on days 0, 4 and 14, whereas Lyt-2.2 mAb on days 0 and 4 in subsequent study.

Direct phenotyping of lymph node cells after




Fig. 1. Clearance of anti-Lyt-2.2 mAb (A) and anti-L3T4 mAb (B) after in vivo administration was examined by antibody-mediated, complement-dependent cytotoxicity assays. Target cells were B6 thymocytes.

in vivo administration of anti-Lyt-2.2 mAb and anti-L3T4 mAb. Change of Lyt-2⁺ cells and L3T4⁺ cells in lymph node after in vivo administration of anti-Lyt-2.2 mAb or anti-L3T4 mAb was then investigated (Fig. 2). Lyt-2⁺ cells




Fig. 2. Phenotypic change of T-cells in lymph node after in vivo administration of anti-Lyt-2.2 mAb (A) and anti-L3T4 mAb (B) was examined by antibody-mediated, complement-dependent cytotoxicity assays, ●, anti-Thy-1.2 mAb; ■, anti-L3T4 mAb; ▲, anti-Lyt-2.2 mAb; ◇, background lysis. N, the reaction of lymph node cells from untreated mice.




Fig. 3. Generation of in vitro cell-mediated cytotoxicity of spleen cells from B6-Lyt-1.1 (O) or B6-Lyt-2.1, 3.1 (Δ) mice that were administered with MEM (control) (A), anti-Lyt-2.2 mAb (B), and anti-L3T4 mAb (C). 3hr ^{51}Cr -release assays were performed at day 23 (A), 22 (B) and 26 (C) after mAb injection. Cytotoxicity was generated by mixed lymphocyte culture in stimulation with irradiated BALB/c spleen cells. Target cells were BALBRL σ^7 1 cells.

were depleted by administration of anti-Lyt-2.2 mAb for about 50 days. L3T4 $^+$ cells were depleted by administration of anti-L3T4 mAb for about 30 days.

Blocking of generation of cytotoxic T-cells from spleen cells of B6 mice administered with anti-Lyt-2.2 mAb. Generation of cytotoxicity from spleen cells after in vitro sensitization with BALB/c spleen cells was investigated with B6Lyt congenic mice that were administered with anti-Lyt-2.2 mAb or anti-L3T4 mAb (Fig. 3). Cytotoxicity was assayed at day 23-26 after administration of mAb. Administration of anti-Lyt-2.2 mAb blocked generation of cytotoxicity from B6-Lyt-1.1 but not B6-Lyt-2.1, 3.1 mice. Administration of anti-L3T4 mAb slightly inhibited generation of cytotoxicity.

Effect of in vivo administration of anti-Lyt-2.2 mAb or anti-L3T4 mAb on tumor rejection. The rejections of a radiation induced leukemia BALBRL σ^7 1 and a RadLV-induced leukemia BALBRVD by CB6F $_1$ recipient mice were studied (Fig. 4). The rejection was blocked by in vivo administration of anti-Lyt-2.2 mAb. Addition of anti-L3T4 mAb to anti-Lyt-2.2 mAb did not alter the results. The rejections were not affect-




Fig. 4. Effect of in vivo administration of anti-Lyt-2.2 mAb (b, f), anti-L3T4 mAb (c, g), both (d, h) or MEM (control) (a, e) on rejection of BALBRL σ^7 1 (a-d) and BALBRVD (e-h) by CB6F $_1$ mice. 5×10^5 BALBRL σ^7 1 and 1×10^6 BALBRVD cells were inoculated into recipient mice.

ed by the administration of anti-L3T4 mAb alone. These results indicated that Lyt-2⁺ cells were fully capable of mediating rejection of these tumors and L3T4⁺ cells appeared not to be involved in rejection. We observed blocking of tumor rejection in mice given only a single injection of Lyt-2 mAb on day 0. Therefore, we investigated the effect of antibody injection at different stages during the course of tumor rejection. As shown in Fig. 5, a single injection of Lyt-2.2 mAb on day 9 after inoculation of B6RV2 was effective for blocking rejection.




Fig. 5. Effect of Lyt-2 mAb injected at different stages during the course of tumor regression. CB6F₁ mice were given a single injection of Lyt-2.2mAb on day 0 (A), or 9 (B) after transplantation of 1×10^6 B6RV2 cells. (C) Untreated control.

DISCUSSION

In the present study, we demonstrated that the rejection of BALBRVD, BALBRL♂1 and B6RV2 tumors in semisyngeneic recipients was blocked by anti-Lyt-2 mAb, but not anti-L3T4 mAb administered in vivo without exogenous complement. It has been widely accepted that cytotoxic T cells are effector cells in tumor rejection. Recently, however, some investigators have raised questions about the functional role of Lyt-2⁺ cells as effectors in rejection of allografts (17) or syngeneic tumors (18), by demonstrating that immunity is passively transferred to immunodeficient mice by spleen cells from which Lyt-2⁺ cells had been eliminated by treatment with anti-Lyt-2 mAb and complement. However, it was also shown that the adult thymectomized, irradiated, and bone marrow-reconstituted mice used in these studies

were not truly T cell deficient (19) and that, in fact, host-derived cytotoxic T cells immune to the allograft were recovered from mice to which Lyt-2⁺-depleted populations were adoptively transferred (20). Thus, it was possible that Lyt-2⁺ cells are involved in graft rejection. Recently (21), it was reported that immunity to syngeneic fibrosarcoma can be adoptively transferred by Lyt-2⁺ cells. Consistent with these findings, immunity is also adoptively transferred by infusion of interleukin 2-dependent cytotoxic T cells to syngeneic tumors (22, 23) and allogeneic tumors (24) or tissue (25). Although our results support these findings and demonstrate directly that Lyt-2⁺ cells are essential for tumor rejection by syngeneic and semisyngeneic recipients, it is still unknown by which mechanism Lyt-2 mAb cause blocking, or at which step of the rejection process Lyt-2 mAb are operative. The finding that a single injection of Lyt-2 mAb on day 0-9 after tumor transplantation effectively blocked rejection suggests that effector cells were functionally blocked rather than that the generation of these cells was inhibited. Previously, we and others demonstrated that in vitro T cell cytotoxicity (26-30) and proliferation (31, 32) in response to alloantigen stimulation, were blocked by anti-Lyt-2 mAb without added complement. We suggested that molecules bearing Lyt-2 and-3 determinants may be involved in T cell recognition. The finding of Lyt-2 blocking of tumor rejection is consistent with these in vitro effects of anti-Lyt-2.

The alteration of the T cell population in the lymph node and spleen of mice injected with mAb is also consistent with findings that, in in vitro cultures of H-2 antigen-stimulated T cells, Lyt-1⁺2⁺3⁺ cells predominate under normal circumstances and Lyt-1⁺2⁻3⁻ cells predominate when Lyt-1⁺2⁺3⁺ cells are blocked by Lyt-2 or-3 antiserum in the absence of complement (31). The decrease of Lyt-2⁺ cells and L3T4⁺ cells in mice injected with Lyt-2 and L3T4 mAb, respectively, did not appear to be due to killing of T cells by mAb and complement from the recipient mice, because Thy-1 and Lyt-1 mAb (both with extremely high titers in cytotoxic tests) did not alter the T cell population.

SUMMARY

Effects of in vivo administration of anti-Lyt-2.2 mAb and anti-L3T4 mAb on the population of T-cells in lymph node, generation of cell-mediated cytotoxicity and tumor rejection were investigated. In vivo administration of anti-Lyt-2.2 mAb selectively eliminated Lyt-2⁺ cells for more than 50 days and that of anti-L3T4 mAb selectively eliminated L3T4⁺ cells for more than 30 days. Generation of cytotoxicity was blocked totally by prior administration of anti-Lyt-2.2 mAb and slightly by anti-L3T4 mAb. By investigating the effects of these mAbs on tumor growth, involvement of Lyt-2⁺ and L3T4⁺ cells for mediating tumor rejections were studied. Rejections of two radiation leukemia virus (RadLV)-induced leukemias, B6RV2 and BALBRVD, and a radiation induced leukemia BALBRL σ^7 1 by CB6F₁ mice were blocked totally by anti-Lyt-2 mAb but not anti-L3T4 mAb. These results indicated that Lyt-2⁺ cells were themselves fully capable of mediating rejection of these tumors. On the other hand, L3T4⁺ cells were not capable of mediating rejection. Injection of anti-Lyt-2.2 mAb on day 9 after tumor inoculation was effective for blocking rejection, suggesting that the effector cells were functionally blocked rather than the generation of those cells was inhibited.

ACKNOWLEDGMENTS

We thank Ms. Yamashita for expert technical assistance and members in Laboratory Animal Center of Nagasaki University for care of mice.

REFERENCES

- 1) Mills, C.D., and R.J. North. 1983. Expression of passively transferred immunity against an established tumor depends on generation of cytolytic T cells in recipients. Inhibition by suppressor T cells. *J. Exp. Med.* 157:1448.
- 2) Rosenstein, M., T.J. Eberlein, and S.A. Rosenberg. 1984. Adoptive immunotherapy of established syngeneic solid tumors: role of T lymphoid subpopulations. *J. Immunol.* 132: 2117.
- 3) Dailey, M.O., E. Pillemer, and I.L. Weissman. 1984. Protection against syngeneic lymphoma by a long-lived cytotoxic T-cell clone. *Proc. Natl. Acad. Sci. USA.* 79: 5384.
- 4) Rosenstein, M., and S.A. Rosenberg. 1984. Generation of lytic and proliferative lymphoid clones to syngeneic tumor: in vitro and in vivo studies. *J. Natl. Cancer Inst.* 72: 1161.
- 5) Eberlein, T.J., M. Rosenstein, and S.A. Rosenberg. 1982. Regression of a disseminated syngeneic solid tumor by systemic transfer of lymphoid cells expanded in interleukin 2. *J. Exp. Med.* 156: 385.
- 6) Engers, H.D., T. Lahaye, G.D. Sorenson, A. L. Glasebrook, C. Horvath, and K.T. Brunner. 1984. Functional activity in vivo of effector T cell populations. II. Anti-tumor activity exhibited by syngeneic anti-MoMuLV-specific cytolytic T cell clones. *J. Immunol.* 133: 1664.
- 7) Fujiwara, H., M. Fukuzawa, T. Yoshioka, H. Nakajima, and T. Hamaoka. 1984. The role of tumor-specific Lyt-1 2 T cells in eradicating tumor cells in vivo. I. Lyt-1 2 T cells do not necessarily require recruitment of host's cytotoxic T cell precursors for implementation of in vivo immunity. *J. Immunol.* 133: 1671.
- 8) Greenberg, P.D., M.A. Cheever, and A. Fefer. 1982. Eradication of disseminated murine leukemia by chemoimmunotherapy with cyclophosphamide and adoptively transferred immune syngeneic Lyt-1 2 lymphocytes. *J. Exp. Med.* 154: 952.
- 9) Nakayama, E., A. Uenaka, E. Stockert, and Y. Obata. 1984. Detection of a unique antigen on radiation leukemia virus-induced leukemia B6RV2. *Cancer Res.* 44: 5138.
- 10) Sato, H., E.A. Boyse, T. Aoki, C. Iritani, and L.J. Old. 1973. Leukemia-associated transplantation antigens related to murine leukemia virus. The X. 1 system: immune response controlled by a locus linked to H-2. *J. Exp. Med.* 138: 96.
- 11) Nakayama, E., H. Shiku, T. Takahashi, H.F. Oettgen, and L.J. Old. 1979. Definition of a unique cell surface antigen of mouse leukemia RL σ^7 1 by cell-mediated cytotoxicity. *Proc. Natl. Acad. Sci. USA.* 76: 3486.
- 12) Dialynas, D.P., Z.S. Quan, K.A. Wall, A. Pierres, J. Quintans, M.R. Loken, M. Pierres and F.W. Fitch. 1983. Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-

- 3/T4 molecule. *J. Immunol.* 131 : 2445.
- 13) Dialynas, D.P., D.B. Wilde, P. Marrack, A. Pierres, K.A. Wall, W. Havran, G. Otten, M. R. Loken, M. Pierres, J. Kappler, and F.W. Fitch. 1983. Characterization of the murine antigenic determinant, designated L 3T 4a, recognized by monoclonal antibody GK1.5 : Expression of L 3T 4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. *Immunol. Rev.* 74 : 29.
 - 14) Ichikawa, T., E. Nakayama, A. Uenaka, M. Monden, and T. Mori. 1987. Effector cells in allelic H-2 class I-incompatible skin graft rejection. *J. Exp. Med.* 166 : 982.
 - 15) Stockert, E., L. J. Old, and E. A. Boyse. 1971. The G_i system. A cell surface alloantigen associated with murine leukemia virus : implication regarding chromosomal integration of the viral genome. *J. Exp. Med.* 133 : 1334.
 - 16) Feeney, A. J., and U. Hammerling. 1976. Ala-1 : a murine alloantigen of activated lymphocytes. I. Serological analysis of mitogen-stimulated T and B cells. *Immunogenetics* 3 : 36.
 - 17) Loveland, B. E., and I. F. C. McKenzie. 1982. Cells mediating graft rejection in the mouse. II. The Ly phenotype of cells producing tumor allograft rejection. *Transplantation (Baltimore)*. 33 : 174.
 - 18) Greenberg, P. D., M. A. Cheever, and A. Fefer. 1981. Eradication of disseminated murine leukemia by chemoimmunotherapy with cyclophosphamide and adoptively transferred immune syngeneic Lyt-1⁺ 2⁻ lymphocytes. *J. Exp. Med.* 154 : 952.
 - 19) Duprez, V., B. Hamilton, and S. J. Burakoff. 1982. Generation of cytolytic T lymphocytes in thymectomized, irradiated, and bone marrow-reconstituted mice. *J. Exp. Med.* 156 : 844.
 - 20) LeFrancois, L., and M. J. Bevan. 1984. A reexamination of the role of Lyt-2-positive T cells in murine skin graft rejection. *J. Exp. Med.* 159 : 57.
 - 21) North, R. J., and I. Bursucker. 1984. Generation and decay of the immune response to a progressive fibrosarcoma I. Ly-1⁺ 2⁻ suppressor T cells down-regulate the generation of Ly-1⁻ 2⁺ effector T cells. *J. Exp. Med.* 159 : 1295.
 - 22) Burton, R. C., and N. L. Warner. 1977. In vitro induction of tumor-specific immunity. IV. Specific adoptive immunotherapy with cytotoxic T cells induced in vitro to plasma-cytoma antigens. *Cancer Immunol. Immunother.* 2 : 91.
 - 23) Eberlein, T. J., M. Rosenstein, and S. A. Rosenberg. 1982. Regression of a disseminated syngeneic solid tumor by systemic transfer of lymphoid cells expanded in interleukin 2. *J. Exp. Med.* 156 : 385.
 - 24) Engers, H. D., A. L. Glasebrook, and G. D. Sorenson. 1982. Allogeneic tumor rejection induced by the intravenous injection of Lyt-2⁺ cytolytic T lymphocyte clones. *J. Exp. Med.* 156 : 1280.
 - 25) Tyler, J. D., S. J. Galli, M. E. Snider, A. M. Dvorak, and D. Steinmuller. 1984. Cloned Lyt-2⁺ cytolytic T lymphocytes destroy allogeneic tissue in vivo. *J. Exp. Med.* 159 : 234.
 - 26) Nakayama, E. H. Shiku, E. Stockert, H. F. Oettgen, and L. J. Old. 1979. Cytotoxic T cells : Lyt phenotype and blocking of killing activity by Lyt antisera. *Proc. Natl. Acad. Sci. USA.* 76 : 1977.
 - 27) Shinohara, N., and D. H. Sachs. 1979. Mouse antibodies capable of blocking cytotoxic T cell function. I. Relationship between the antigen reactive with blocking antibodies and the Lyt-2 locus. *J. Exp. Med.* 150 : 432.
 - 28) Hollander, N., E. Pillemer, and I. L. Weissman. 1980. Blocking effect of Lyt-2 antibodies on T cell functions. *J. Exp. Med.* 152 : 674.
 - 29) Sarmiento, M., A. L. Glasebrook, and F. W. Fitch. 1980. IgG or IgM monoclonal antibodies reactive with different determinants on the molecular complex bearing Lyt-2 antigen block T cell-mediated cytotoxicity in the absence of complement. *J. Immunol.* 125 : 2665.
 - 30) Fan, J., and B. Bonavida. 1981. Studies on the induction and expression of T cell-mediated immunity. XII. The concomitant loss and recovery of membrane-associated Lyt-2 antigens, lymphocyte-target cell binding, and the antigen-specific and-nonspecific cytotoxic activity of alloimmune T lymphocytes after treatment with trypsin. *J. Immunol.* 127 : 1856.
 - 31) Nakayama, E., W. Dippold, H. Shiku, H. F. Oettgen, and L. J. Old. 1980. Alloantigen-induced T-cell proliferation : Lyt phenotype of responding cells and blocking of proliferation by Lyt antisera. *Proc. Natl. Acad. Sci. USA.* 77 : 2890.
 - 32) Hollander, N., E. Pillemer, and I. L. Weiss-

man, 1981. Effect of Lyt antibodies on T-cell functions : augmentation by anti-Lyt- 1 as opposed to inhibition by anti-Lyt-2. Proc. Natl. Acad. Sci. USA. 78 : 1148.