



Title	Dengue virus, Nepal
Author(s)	Pandey, Basu Dev; Morita, Kouichi; Khanal, Santa Raj; Takasaki, Tomohiko; Miyazaki, Isao; Ogawa, Tetsuro; Inoue, Shingo; Kurane, Ichiro
Citation	Emerging Infectious Diseases, 14(3), pp.514-515; 2008
Issue Date	2008-03
URL	http://hdl.handle.net/10069/18710
Right	

This document is downloaded at: 2019-03-20T15:56:47Z

10. Winger DA, Fass RJ. Antibiotic cement and beads for orthopedic infections. *Antimicrob Agents Chemother.* 1996;40:2675-9.

Address for correspondence: Jason J. Schafer, UPMC St. Margaret, Department of Pharmacy, 815 Freeport Rd, Pittsburgh, PA 15215, USA; email: schaferrjj@upmc.edu

Dengue Virus, Nepal

To the Editor: *Dengue virus* belongs to the genus *Flavivirus*, family *Flaviviridae*. It has 4 serotypes: dengue virus type 1 (DENV-1), dengue virus type 2 (DENV-2), dengue virus type 3 (DENV-3), and dengue virus type 4 (DENV-4). Dengue virus is maintained in a cycle between humans

and *Aedes aegypti*, domestic day-biting mosquitoes. Dengue virus induces clinical illness, which ranges from a nonspecific viral syndrome (dengue fever [DF]) to severe and fatal hemorrhagic disease (dengue hemorrhagic fever [DHF]). DF/DHF occurs primarily in tropical and subtropical areas of the world. Domestic dengue virus infection occurs in >100 countries; >2.5 billion persons live in these areas. Approximately 100 million cases of DF, 500,000 cases of DHF, and several thousand deaths occur annually worldwide (1). During the past decades, dengue virus has emerged in southern Asia; DF/DHF epidemics have occurred in Bhutan, India, Maldives, Bangladesh, and Pakistan (2-4).

From August through November 2006, the number of febrile patients increased in 4 major hospitals in the Terai region of Nepal: Nepalgunj Medical College, Bheri Zonal Hospital in Nepalgunj, Tribhuban Hospital in Dang, and Narayani subregional

hospital in Birgunj. Patients with severe symptoms were referred to Sukraj Tropical and Infectious Disease Hospital, Kathmandu, for diagnosis and treatment. The clinical features in most patients were consistent with signs of DF, but some patients showed signs (high fever, rash, ecchymosis, epistaxis, positive tourniquet test, liver dysfunction, and thrombocytopenia [platelet count <100,000/mm³]) consistent with the World Health Organization (WHO) definition of DHF. Ascites and plural effusion developed in 2 patients. Blood specimens were collected from all patients at the time of admission to the local hospitals. Particle agglutination (PA) assay (Pentax Ltd, Tokyo, Japan) (5) and immunoglobulin (Ig) M-capture ELISA (Dengue/JE IgM Combo ELISA kit, Panbio Ltd, Brisbane, Queensland, Australia) were performed. Dengue virus-specific IgM was detected in 11 patients who had fever, headache, and rash (Table). Each of these patients had negative

Table. Clinical and laboratory data for 11 patients admitted to hospitals and diagnosed with dengue fever or dengue hemorrhagic fever, Nepal, 2006*

Patient age, y/Sex	Month admitted	Location	Initial diagnosis	Travel history	Clinical signs and symptoms	Selected laboratory and other test results
20/M	Sep	Kathmandu	DF	Yes	Fever, headache, nausea	Hb 15.4 g/dL; TLC 10,500/mm ³ ; Plt 185,000/mm ³ ; blood culture for salmonellae negative; ALT 38 IU/L
27/F	Sep	Bardiya	Viral fever	No	Fever, headache, vomiting	TLC 5,600/mm ³ ; blood culture for salmonellae negative
3/M	Sep	Salayan	Encephalitis	No	Fever, vomiting, convulsions	Widal negative; TLC 4,700/mm ³
13/M	Oct	Sindhuli	Typhoid fever	No	Fever, headache	Widal negative; TLC 4,500/mm ³ ; blood culture for salmonellae negative; <i>Brucella</i> antigen negative; chest radiograph normal
22/M	Oct	Birgunj	DHF	No	Fever, headache, vomiting, ascites	Bil 0.8 mg/dL; ALT 80 IU/L; Plt 22,000/mm ³ ; chest radiograph normal
55/F	Oct	Dang	DF	No	Fever, headache, muscular pain	Plt 51,000/mm ³ ; TLC 7,600/mm ³ ; MP negative; ESR 20 mm/h; Bil 0.7 mg/dL
22/F	Oct	Birgunj	Viral fever	No	Fever, headache, body ache	<i>Brucella</i> negative; Widal negative; TLC 5,600/mm ³
13/M	Nov	Dang	DF	No	Fever, headache, rashes	Plt 95,000/mm ³ ; TLC 4,700/mm ³ ; Hb 13.1 g%; Bil 0.8mg/dL; ALT 26 IU/L
35/F	Nov	Birgunj	DHF	No	Fever, headache, bruises; tourniquet: positive	Bil 0.81mg/dL; Plt 31,000/mm ³ ; PT 2 min 30 s (control 14)
40/M	Nov	Birgunj	DF	No	Fever, headache, rashes	ALT 127IU/L; Plt 110,000 /mm ³ ; PCV 38.8%;TLC 5,500/mm ³ ; ultrasonography liver size, 16.8 cm
42/M	Nov	Dang	DF	No	Fever, headache, rashes	Bil 0.7 mg/dL; Widal test negative; TLC 6,800/mm ³ ; Plt 164,000/mm ³

*Blood specimens were collected at time of hospital admission. Diagnosis was confirmed by using immunoglobulin M-capture ELISA. DF, dengue fever; Hb, hemoglobin; TLC, total leukocyte count; Plt, platelets; ALT, alanine aminotransferase; DHF, dengue hemorrhagic fever; Bil, bilirubin; MP, malaria parasites; ESR, erythrocyte sedimentation rate; PT, prothrombin time; PCV, packed cell volume.

results for Japanese encephalitis virus-specific IgM. Of the 11 patients, 10 had no history of travel to India or other dengue-endemic countries. DF or DHF was initially diagnosed for 7 patients, and viral encephalitis, typhoid fever, or viral fever was diagnosed for others without serologic tests. Reverse transcription-PCR and virus isolation were performed at Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan, but the dengue virus genome was not detected, and no virus was isolated, likely because sample collection was delayed and the sample was transported to Japan in a deteriorated condition.

DF/DHF have been considered to be a possible public health threat to Nepal because DF/DHF epidemics have occurred recently in India and Pakistan, which reported several thousand cases and >100 deaths (6). The first DF case in Nepal was reported in 2004 (7). Further, the first DENV-2 strain of Nepal origin was isolated from a Japanese traveler who visited Nepal and in which DF developed after the patient returned to Japan. The isolated DENV-2 (GenBank accession no. AB194882) was 98% homologous with DENV-2 isolated in India (8). The prevalence of dengue virus antibody was reported to be 10.4% in the southwestern region of Nepal (9). These reports suggest that dengue virus has been circulating in Nepal for several years. Thus, DF/DHF has likely been misdiagnosed and illness caused by dengue virus underestimated in Nepal. In contrast, Japanese encephalitis has been a public health problem in southwestern region of Nepal, and large epidemics have occurred almost every year since 1978 (10). Nepal has no dengue surveillance programs, and health professionals do not usually consider dengue as a differential diagnosis.

The emergence occurred in the lowland Terai belt region, which borders the state of Bihar, India. The *Aedes* mosquito is known to persist in

this region. The emerging DENV-2 is likely to have been introduced into Nepal from India.

**Basu Dev Pandey,*
Kouichi Morita,†
Santa Raj Khanal,‡
Tomohiko Takasaki,§
Isao Miyazaki,¶
Tetsuro Ogawa,¶
Shingo Inoue,†
and Ichiro Kurane§**

*Tropical and Infectious Disease Hospital, Kathmandu, Nepal; †Nagasaki University, Nagasaki, Japan; ‡Tribhuban University, Kathmandu, Nepal; §National Institute of Infectious Diseases, Tokyo, Japan; and ¶Pentax Company Limited, Tokyo, Japan

References

1. World Health Organization. Dengue and dengue haemorrhagic fever. Fact sheet no.117; revised 2002 Apr [cited 2006 Oct 31]. Available from <http://www.who.int/mediacentre/factsheets/fs117/en>
2. World Health Organization. Dengue outbreak in Bhutan. Communicable Disease Newsletter. 2007;4(1).
3. Islam MA, Ahmed M, Begum N, Choudhury N, Khan A, Parquet C, et al. Molecular characterization and clinical evaluation of dengue outbreak in 2002 in Bangladesh. *Jpn J Infect Dis.* 2006;59:85–91.
4. Jamil B, Hasan R, Zafar A, Bewley K, Chamberlain J, Mioulet V, et al. Dengue virus serotype, Karachi, Pakistan. *Emerg Infect Dis.* 2007;13:182–3.
5. Pandey B, Yamamoto A, Morita K, Kurosawa Y, Rai S, Adhikari S, et al. Serodiagnosis of Japanese encephalitis among Nepalese patients by the particle agglutination assay. *Epidemiol Infect.* 2003;131:881–5.
6. Gupta E, Dar L, Kapoor G, Broor S. The changing epidemiology of dengue in Delhi. *Virology.* 2006;3:92.
7. Pandey BD, Rai SK, Morita K, Kurane I. First case of dengue in Nepal. *Nepal Med Coll J.* 2004;6:157–9.
8. Takasaki T, Kotaki A, Nishimura K, Sato Y, Tokuda A, Lim C, et al. Dengue virus type 2 from an imported dengue patient in Japan. First isolation of dengue virus from Nepal. *J Travel Med.* 2007;14:445–8.
9. Sherchand JB, Pandey BD, Haruki K, Jimba M. Sero-diagnosis of Japanese encephalitis and dengue virus infection from clinically suspected patients of Nepal. *J Inst Med.* 2001;23:25–31.
10. Epidemiology and Disease Control Division, Ministry of Health Nepal. The annual report of malaria, Kala-azar and Japanese encephalitis in Nepal, 2005. Kathmandu, Nepal: Ministry of Health; 2006.

Address for correspondence: Basu Dev Pandey, Tropical and Infectious Disease Hospital, Teku Road, GPO Box 9045, Kathmandu, Nepal; email: basupandey@wlink.com.np

Human Tuberculosis Caused by *Mycobacterium bovis*, Taiwan

To the Editor: *Mycobacterium bovis* is one of the causative agents of tuberculosis (TB) in humans and animals. Drinking unpasteurized milk, eating undercooked meat, and close contact with infected animals are the main sources of infection for humans. Currently, 119 *M. bovis* spoligotypes are contained in the fourth international spoligotyping database (SpolDB 4) and are categorized into 3 main sublineages corresponding to ST prototypes 482, 683, and 479 (1).

Although an *M. bovis* surveillance program for farm animals has been implemented by the Taiwan Council of Agriculture, no surveillance system exists for human TB cases caused by *M. bovis*. To monitor the epidemiology of *M. bovis* in domestic animals, a regular tuberculin skin test (TST) is compulsory for cattle and sheep and optional for deer in Taiwan (2). In 2005, screening of *Mycobacterium* spp. infections by TST was performed for 111,412 cattle and 73,396 caprine and ovine herds, of which 188 (0.17%) and 148 (0.2%), respectively, were positive (2). We used spacer oligonucleotide typing (spoligotyping) and