



Title	Homeomorphism Groups of Homogeneous Spaces
Author(s)	Karube, Takashi
Citation	長崎大学教育学部自然科学研究報告. vol.33, p.5-9; 1982
Issue Date	1982-02-28
URL	http://hdl.handle.net/10069/32581
Right	

This document is downloaded at: 2018-11-18T04:14:26Z

Homeomorphism Groups of Homogeneous Spaces

Takashi KARUBE

Department of Mathematics, Faculty of Education
Nagasaki University, Nagasaki
(Received Oct. 31, 1981)

Abstract

Let X be a separable metrizable coset-space of a locally compact group, which has a local cross-section and admits a nontrivial flow. Let $\mathcal{H}(X)$ be the group of homeomorphisms on X , endowed with the compact-open topology, and $\mathcal{H}(X, x)$ the subspace of $\mathcal{H}(X)$ consisting of those homeomorphisms which fix a point x of X . Then $\mathcal{H}(X)$ is an l_2 -manifold if and only if X is an ANR and $\mathcal{H}(X, x)$ is an l_2 -manifold. Among applications of this, we see that if X is the plane \mathbb{R}^2 , or punctured real projective plane, or punctured torus, then $\mathcal{H}(X)$ is an l_2 -manifold.

Introduction

Let $\mathcal{H}(X)$ be the group of homeomorphisms on a topological space X , endowed with the compact-open topology. For twelve years now there has been considerable interest in the question of whether $\mathcal{H}(X)$ is an l_2 -manifold for a manifold X . Among (finite-dimensional) manifolds for which the question is affirmatively answered there are metrizable connected 1-manifolds (R. D. Anderson [1], T. Karube [5]), compact metric 2-manifolds (R. Luke and W. K. Mason [7], H. Toruńczyk [9]). And it is reported that if X is the plane \mathbb{R}^2 less an infinite closed discrete set then $\mathcal{H}(X)$ is not locally contractible (J. Keesling [6], p. 2). In this paper, for such a separable metrizable coset-space X of a locally compact group that has a local cross-section and admits a nontrivial flow, it is shown that $\mathcal{H}(X)$ is an l_2 -manifold if and only if X is an ANR and $\mathcal{H}(X, x)$ is an l_2 -manifold, where $\mathcal{H}(X, x)$ is the subspace of $\mathcal{H}(X)$ consisting of those homeomorphisms which fix a point x of X . Joining this fact and a result of R. Arens, we see that if X is a positive-dimensional compact coset-space of a Lie group then to be an l_2 -manifold for $\mathcal{H}(X)$ and $\mathcal{H}(X-x)$ is equivalent ($x \in X$). Applying the results of R. Luke, W. K. Mason and H. Toruńczyk to this, we know that if X is a compact

2-manifold homeomorphic to a coset-space of a Lie group then $\mathcal{H}(X-x)$ is an l_2 -manifold ($x \in X$). Thus we obtain non-compact 2-manifolds whose homeomorphism groups are l_2 -manifolds.

1. Homeomorphism groups of locally compact homogeneous spaces.

Let G be a locally compact Hausdorff topological group, $X = G/H$ the left coset-space of G by a closed subgroup H . Assume that the coset-space X has a local cross-section. Let $\mathcal{H}(X)$ (resp. $\mathcal{H}(X-x)$) be the group of homeomorphisms on X (resp. $X-x$, where $x \in X$), endowed with the compact-open topology, and $\mathcal{H}(X, x)$ the subspace of $\mathcal{H}(X)$ consisting of those homeomorphisms which fix a point $x \in X$. These notations $G, X, \mathcal{H}(X), \mathcal{H}(X, x), \mathcal{H}(X-x)$ will keep these meanings throughout the paper.

THEOREM 1. *Let $G, X, \mathcal{H}(X), \mathcal{H}(X, x)$ be those as above. Let x_0 be the coset containing the identity of G , and p the projection of $\mathcal{H}(X)$ onto X defined by $p(\phi) = \phi(x_0)$ ($\phi \in \mathcal{H}(X)$). Then for each $x \in X$ there exists a neighborhood U of x in X such that $p^{-1}(U) \approx U \times \mathcal{H}(X, x_0)$.*

(Here “ \approx ” means “homeomorphic to”, and “ $A \times B$ ” means “the product space of spaces A and B .”)

PROOF. As the space X is locally compact Hausdorff, the product operation in $\mathcal{H}(X)$ is continuous and the translations in $\mathcal{H}(X)$ are homeomorphisms, while the inverse operation in $\mathcal{H}(X)$ is not necessarily continuous. Let π be the natural projection of G onto X , l_a the left translation in X by $a \in G$ defined by $l_a(x) = \pi(ab)$ ($b \in x \in X$), and $\mathcal{L} = \{l_a | a \in G\}$. \mathcal{L} is a subgroup of $\mathcal{H}(X)$. Let ω be the mapping $G \rightarrow \mathcal{L}$ defined by $\omega(a) = l_a$ ($a \in G$), then ω is a continuous surjective homomorphism. For an arbitrary point $x \in X$, let f be a local cross-section defined on an open neighborhood U of x and put $W = (\omega \circ f)(U)$. Then $p|W$ and $\omega \circ f$ are homeomorphisms $W \rightarrow U$, $U \rightarrow W$ respectively and they are inverses of each other. On the other hand let \mathcal{H}^* be the coset-space $\mathcal{H}(X) / \mathcal{H}(X, x_0)$, π^* the natural projection $\mathcal{H}(X) \rightarrow \mathcal{H}^*$, and let $q = p \circ \pi^{*-1}$. The mapping q is a continuous bijection $\mathcal{H}^* \rightarrow X$ and $\pi^*|W$ is a continuous bijection $W \rightarrow q^{-1}(U)$. Thus $p^{-1}(U) = W \circ \mathcal{H}(X, x_0)$. Now let Φ be the mapping $W \times \mathcal{H}(X, x_0) \rightarrow p^{-1}(U)$ defined by $\Phi((w, \phi)) = w \circ \phi$ ($(w, \phi) \in W \times \mathcal{H}(X, x_0)$), then Φ is a continuous bijection. To show that Φ^{-1} is continuous, put $N = \ker \omega$, and let G' be the factor group G/N , π' the natural projection $G \rightarrow G'$, and let $x' = \pi' \circ f(x)$. We can choose a closure compact, open symmetric neighborhood V' of x' in G' and an open neighborhood U_1 of x in X such that $f(U_1) \subset \pi'^{-1}(V')$ and $U_1 \subset U$. Put $W_1 = (\omega \circ f)(U_1)$. The mapping $\omega' = \omega \circ \pi'^{-1}: G' \rightarrow \mathcal{L}$ is a continuous isomorphism and its restriction on $\text{Cl } V'$ is a homeomorphism such that $W_1 \subset \omega'(\text{Cl } V')$. Since $\omega'(\text{Cl } V')$ is symmetric, the mapping $W_1 \rightarrow \omega'(\text{Cl } V')$ which maps w to w^{-1} ($w \in W_1$) is well-defined

and continuous. Using the fact we can prove that $\Phi^{-1}|_{W_1 \circ \mathcal{H}(X, x_0)}$ is continuous. Consequently

$$p^{-1}(U_1) = W_1 \circ \mathcal{H}(X, x_0) \approx U_1 \times \mathcal{H}(X, x_0).$$

COROLLARY (J. Keesling). *If G is a locally compact Hausdorff topological group, then $\mathcal{H}(G)$ is a product space over G .*

PROOF. In this case we can consider in the proof of Theorem 1 that $H = \{\text{the identity}\}$, $X = G = U$, $W = \mathcal{L} \approx G$, and \mathcal{L} is a topological group.

The fact in Corollary is found in a remark of J. Keesling [6, p.15].

Now from Theorem 1 we see that $\mathcal{H}(X)$ is locally homeomorphic to the product space $X \times \mathcal{H}(X, x)$. Thus in particular for $\mathcal{H}(X)$ to be locally contractible (or locally connected), X must be so.

A space is called an l_2 -manifold if it is separable metrizable space which is locally homeomorphic to l_2 .

On the condition for $\mathcal{H}(X)$ to be an l_2 -manifold we have the following.

THEOREM 2. *Let X be a separable metrizable coset-space of a locally compact Hausdorff topological group. Assume that X has a local cross-section and admits a nontrivial flow. Then $\mathcal{H}(X)$ is an l_2 -manifold if and only if X is an ANR and $\mathcal{H}(X, x)$ is an l_2 -manifold for a fixed point $x \in X$.*

(Here "ANR" means "absolute neighborhood retract for the class of all metrizable spaces.")

PROOF. Since X is separable metrizable, locally compact Hausdorff, both $\mathcal{H}(X)$ and $\mathcal{H}(X, x)$ are separable metrizable. If $\mathcal{H}(X)$ is an l_2 -manifold, then $X \times \mathcal{H}(X, x)$ is an l_2 -manifold and so a local ANR, hence it is an ANR. Then both X and $\mathcal{H}(X, x)$ are ANR's. By a theorem of Toruńczyk [9], $\mathcal{H}(X, x)$ is an l_2 -manifold. The converse can be easily proved using the properties of ANR's and the theorem of Toruńczyk.

2. Homeomorphism groups of locally connected, compact homogeneous spaces.

We apply Theorem 2 to $\mathcal{H}(X)$ for locally connected compact coset-spaces X , and show a sufficient condition for $\mathcal{H}(X)$ and $\mathcal{H}(X-x)$ to be l_2 -manifolds is equivalent, while taking account of the assumption "local connectedness", propositions hereafter can be derived basing on a McCarty's result [8, p. 295] also.

LEMMA (R. Arens). *Let X be a locally connected, compact Hausdorff space, and x an arbitrary point of X . Then $\mathcal{H}(X, x)$ is topologically isomorphic to $\mathcal{H}(X-x)$.*

PROOF. As McCarty [8] has remarked, this is shown by Theorems 1 3, and 4 of [3], and Theorem 2 of [2].

By Theorem 2 and Lemma we have the following.

THEOREM 3. *Let X be a locally connected, compact metrizable coset-space of a locally compact Hausdorff topological group. Assume that X is an ANR and has a local cross-section and admits a nontrivial flow. Then $\mathcal{H}(X)$ is an l_2 -manifold if and only if $\mathcal{H}(X-x)$ is an l_2 -manifold where x is any fixed point of X .*

REMARK 1. A coset-space G/H of a locally compact Hausdorff topological group G has a local cross-section if 1) G is separable metrizable finite-dimensional, or 2) there are no arbitrarily small nontrivial subgroups in H , or 3) all arbitrarily small subgroups of G are in H (T. Karube [4]).

REMARK 2. Following space admit a nontrivial flow : 1) a metric space that contains a subset Y with nonempty interior such that Y is homeomorphic to \mathbb{R}^n or l_2 or I^∞ , 2) a locally compact Hausdorff topological group which is not totally disconnected (J. Keesling [6]).

COROLLARY 1. *If X is a positive-dimensional compact coset-space of a Lie group, then the same conclusion as in Theorem 3 holds.*

EXAMPLES. The following spaces satisfy the condition on X in Corollary 1 : torus T^n , sphere S^n , projective space RP_n , CP_n , or HP_n over the field of real numbers, complex numbers, or quaternions respectively ($n > 0$) — $S^n \approx O(n+1)/O(n)$, $RP_n \approx O(n+1)/(O(n) \times S^0)$, $CP_n \approx U(n+1)/(U(n) \times S^1)$, $HP_n \approx Sp(n+1)/(Sp(n) \times S^3)$. ($O(n)$: the orthogonal group, $U(n)$: the unitary group, $Sp(n)$: the symplectic group)

COROLLARY 2. *If X is a compact 2-manifold homeomorphic to a coset-space of a Lie group, then $\mathcal{H}(X-x)$ is an l_2 -manifold for each point x of X .*

PROOF. By a theorem of Toruńczyk [9], for a separable metric space M which admits a nontrivial flow, $\mathcal{H}(M)$ is an l_2 -manifold if and only if it is an ANR. When M is a compact metric 2-manifold without boundary, Luke and Mason [7] have shown that $\mathcal{H}(M)$ is an ANR. Hence by Corollary 1 we see that $\mathcal{H}(X-x)$ is an l_2 -manifold.

EXAMPLES. The following spaces satisfy the condition on $X-x$ in Corollary 2 : plane \mathbb{R}^2 , punctured torus, punctured real projective plane, etc.

REMARK. We can treat analogously "local contractibility" in place of " l_2 -manifold". Therefore if X is a compact coset-space of a Lie group, then $\mathcal{H}(X-x)$ is locally contractible, by a theorem of A. V. Černavskii (Math. USSR Sb. 8 (1969)). Thus, for example, if Y is a Euclidean space, or a punctured torus, or a punctured projective space over the field of real numbers, complex numbers, or quaternions, then $\mathcal{H}(Y)$ is locally contractible.

References

- [1] R. D. Anderson : *Spaces of homeomorphisms of finite graphs*. (Manuscript)
- [2] R. Arens : *A topology for spaces of transformations*, Ann. of Math. (2) **47** (1946) 480-495.
- [3] R. Arens : *Topologies for homeomorphism groups*, Amer. J. Math. **68** (1946), 593-610.
- [4] T. Karube : *On the local cross-sections in locally compact groups*, J. Math. Soc. Japan **10** (1958), 343-347.
- [5] T. Karube : *Topological description of the space of homeomorphisms on connected 1-manifolds*. To appear.
- [6] J. Keesling : *Using flows to construct Hilbert space factors of function spaces*, Trans. Amer. Math. Soc. **161** (1971), 1-24.
- [7] R. Luke and W. K. Mason : *The space of homeomorphisms on a compact two-manifold is an absolute neighborhood retract*, Trans. Amer. Math. Soc. **164** (1972), 275-285.
- [8] G. S. McCarty : *Homeotopy groups*, Trans. Amer. Math. Soc. **106** (1963), 293-304.
- [9] H. Toruńczyk : *Absolute retracts as factors of normed linear spaces*, Fund. Math. **86** (1974), 53-67.